

3

FILE 'HOME' ENTERED AT 11:57:42 ON 29 NOV 2000

=> file caplus medline embase wpids biosis

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.15	0.15

FILE 'CAPLUS' ENTERED AT 11:58:21 ON 29 NOV 2000
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2000 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'MEDLINE' ENTERED AT 11:58:21 ON 29 NOV 2000

FILE 'EMBASE' ENTERED AT 11:58:21 ON 29 NOV 2000
COPYRIGHT (C) 2000 Elsevier Science B.V. All rights reserved.

FILE 'WPIDS' ENTERED AT 11:58:21 ON 29 NOV 2000
COPYRIGHT (C) 2000 DERWENT INFORMATION LTD

FILE 'BIOSIS' ENTERED AT 11:58:21 ON 29 NOV 2000
COPYRIGHT (C) 2000 BIOSIS(R)

=> s phytostenol# or phytosterol# or sitostenol# or sitostanol# or sitosterol#

L1 18440 PHYTOSTENOL# OR PHYTOSTEROL# OR SITOSTENOL# OR SITOSTANOL# OR SITOSTEROL#

=> s fatty(w)acid#

L2 535195 FATTY(W) ACID#

=> s unsaturat?(w)fatty(w)acid#

L3 35497 UNSATURAT?(W) FATTY(W) ACID#

=> s l1 and l3

L4 225 L1 AND L3

=> s hypocholest? or lower(n)cholest? or reduct?(s)cholest?

L5 53398 HYPOCHOLEST? OR LOWER(N) CHOLEST? OR REDUCT?(S) CHOLEST?

=> s l4 and l5

L6 11 L4 AND L5

=> d kwic ibib so l6 1-11

L6 ANSWER 1 OF 11 CAPLUS COPYRIGHT 2000 ACS

TI Method for manufacturing fat-soluble **phytosterol** or phytostanol ester of **unsaturated fatty acid**

AB The present invention provides a method for manufg. fat-sol. **phytosterol** or phytostanol ester of **unsatd. fatty acid** for inhibiting the absorption of cholesterol and foodstuffs contg. the same. The method for manufg. fat-sol.

phytosterol or phytostanol ester of **unsatd. fatty acid** comprises the steps of: esterification of **phytosterol** or phytostanol with **unsatd. fatty acid** by dissolving them in a nonpolar org. solvent with a basic catalyst and adding a carboxyl group activating agent dissolved.

ST **phytosterol** fatty ester fat soluble prodn **cholesterol** absorption **redn**; phytostanol fatty ester fat soluble prodn **cholesterol** absorption **redn**; **sitosterol** oleate prodn esterification **cholesterol** lowering effect

IT Fatty acids, preparation
 RL: BUU (Biological use, unclassified); IMF (Industrial manufacture);

BIOL (Biological study); PREP (Preparation); USES (Uses)
 (esters; method for manufg. fat-sol. **phytosterol** or phytostanol ester of **unsatd. fatty acid**)

IT 57-88-5, **Cholesterol**, biological studies
 RL: BPR (Biological process); BIOL (Biological study); PROC (Process)
 (method for manufg. fat-sol. **phytosterol** or phytostanol ester of **unsatd. fatty acid**)

IT 3712-16-1P, .beta.-**Sitosterol** oleate
 RL: BUU (Biological use, unclassified); IMF (Industrial manufacture);

BIOL (Biological study); PREP (Preparation); USES (Uses)
 (method for manufg. fat-sol. **phytosterol** or phytostanol ester of **unsatd. fatty acid**)

IT 83-46-5, .beta.-**Sitosterol** 112-80-1, Oleic acid, reactions
 RL: RCT (Reactant)
 (method for manufg. fat-sol. **phytosterol** or phytostanol ester of **unsatd. fatty acid**)

ACCESSION NUMBER: 2000:742194 CAPLUS
 DOCUMENT NUMBER: 133:323278
 TITLE: Method for manufacturing fat-soluble **phytosterol** or phytostanol ester of **unsaturated fatty acid**

INVENTOR(S): Chung, Dae-Won; Noh, Seung-Kwon; Kim, Kab-Sig
 PATENT ASSIGNEE(S): Eugene Science, Inc., S. Korea
 SOURCE: PCT Int. Appl., 19 pp.
 CODEN: PIXXD2

DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000061694	A1	20001019	WO 1999-KR569	19990921
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
PRIORITY APPLN. INFO.:			KR 1999-12965	19990413
SO PCT Int. Appl., 19 pp. CODEN: PIXXD2				
REFERENCE COUNT:		4		
REFERENCE(S):		(1) Henkel; WO 9925362 A1 1999 CAPLUS (2) Raisio Benecol Oy; WO 9956558 A1 1999 CAPLUS (3) Raisio Margarini; WO 9219640 A1 1992 CAPLUS (4) Tackett, T; US 5117016 A 1992 CAPLUS		

AB . . . plasma cholesterol level has a direct assocn. with coronary heart

disease, ischemic heart disease, acute myocardial infarction, and atherosclerosis. A **lower cholesterol** level reduces their risk. Among the plasma cholesterol fractions the low-d.

lipoprotein

cholesterol and first of all its oxidized forms have strong relationship to the mentioned diseases. The chem. similar **phytosterols** in plants have favorable effect, they prevent the absorption of cholesterol from the intestine. The high-d. lipoprotein transports the cholesterol. . . may be influenced by nutrition. It looks advantageous the diet contg. low fat, low satd. fatty acids, low cholesterol, more **unsatd. fatty acids**, high consumption of fruits and vegetables, adequate intake of dietary fiber. The high-d. lipoprotein level will be increased mainly. . .

ACCESSION NUMBER: 1999:521301 CAPLUS
DOCUMENT NUMBER: 131:228043
TITLE: The essential and accursed cholesterol
AUTHOR(S): Biro, Gyorgy
CORPORATE SOURCE: Budapest, Hung.
SOURCE: Elelmez. Ip. (1999), 53(6), 161-166
CODEN: EMIPAB; ISSN: 0013-5909
PUBLISHER: METE
DOCUMENT TYPE: Journal; General Review
LANGUAGE: Hungarian
SO Elelmez. Ip. (1999), 53(6), 161-166
CODEN: EMIPAB; ISSN: 0013-5909

L6 ANSWER 3 OF 11 CAPLUS COPYRIGHT 2000 ACS

TI Use of mixtures containing **phytostenols** for producing **hypocholesteremic** preparations

AB Mixts. of active agents contg. (a) **phytostenols** and/or **phytostenol** esters and (b) conjugated fatty acids or their glycerides are used to produce **hypocholesteremic** prepns. These mixts. have a synergistic effect in reducing the cholesterol content of serum. When encapsulated in gelatin, the prepns.. . .

ST **hypocholesteremic phytostenol unsatd fatty acid**; synergistic **hypocholesteremic phytostenol fatty acid**

IT **Unsaturated fatty acids**

RL: BAC (Biological activity or effector, except adverse); FFD (Food or feed use); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (diunsatd., with conjugated double bonds; use of mixts. contg. **phytostenols** for producing **hypocholesteremic** prepns.)

IT Sterol esters

Sterols

RL: BAC (Biological activity or effector, except adverse); FFD (Food or feed use); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (from plants; use of mixts. contg. **phytostenols** for producing **hypocholesteremic** prepns.)

IT Glycerides, biological studies

RL: BAC (Biological activity or effector, except adverse); FFD (Food or feed use); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (polyunsatd. fatty acid-contg., with conjugated double bonds; use of mixts. contg. **phytostenols** for producing **hypocholesteremic** prepns.)

IT Anticholesteremic agents

Butter

Capsules (drug delivery systems)

Cocoa products

Dietary food

Food

Margarine

Mayonnaise

Salad dressings

Sausage
 Synergistic drug interactions
 (use of mixts. contg. **phytostenols** for producing
hypocholesteremic prepns.)

IT Fats and Glyceridic oils, biological studies
 RL: FFD (Food or feed use); BIOL (Biological study); USES (Uses)
 (use of mixts. contg. **phytostenols** for producing
hypocholesteremic prepns.)

IT Polyunsaturated fatty acids
 RL: BAC (Biological activity or effector, except adverse); FFD (Food or
 feed use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (with conjugated double bonds; use of mixts. contg.
phytostenols for producing **hypocholesteremic** prepns.)

IT Fatty acid esters
 RL: BAC (Biological activity or effector, except adverse); FFD (Food or
 feed use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (with **phytostenols**; use of mixts. contg. **phytostenols**
 for producing **hypocholesteremic** prepns.)

IT 83-45-4, .beta.-**Sitostanol** 83-45-4D, .beta.-**Sitostanol**
 , esters 83-46-5 83-46-5D, esters 1839-11-8D, 9,11-Octadecadienoic
 acid, esters with **phytostenols** 41005-65-6 109033-78-5
 RL: BAC (Biological activity or effector, except adverse); FFD (Food or
 feed use); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (use of mixts. contg. **phytostenols** for producing
hypocholesteremic prepns.)

ACCESSION NUMBER: 1999:344854 CAPLUS
 DOCUMENT NUMBER: 130:347399
 TITLE: Use of mixtures containing **phytostenols** for
 producing **hypocholesteremic** preparations
 INVENTOR(S): Fabry, Bernd
 PATENT ASSIGNEE(S): Henkel Kommanditgesellschaft auf Aktien, Germany
 SOURCE: PCT Int. Appl., 19 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9925362	A1	19990527	WO 1998-EP7059	19981105
W: AU, BG, BR, BY, CA, CN, CZ, HU, ID, IS, JP, KR, LT, LV, MX, NO, NZ, PL, RO, RU, SI, SK, TR, UA, US RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
DE 19750453	A1	19990527	DE 1997-19750453	19971114
AU 9915603	A1	19990607	AU 1999-15603	19981105
EP 1028733	A1	20000823	EP 1998-959848	19981105
R: DE, ES, FR, GB, IT, NL				
PRIORITY APPLN. INFO.:			DE 1997-19750453	19971114
			WO 1998-EP7059	19981105
OTHER SOURCE(S): MARPAT 130:347399				
SO PCT Int. Appl., 19 pp.				
CODEN: PIXXD2				
REFERENCE COUNT: 7				
REFERENCE(S):				
(1) Funes; 1980, 5, CAPLUS				
(2) Funes, C; AN ASOC QUIM ARGENT 1978, V66(5), P239				
(3) Hasegawa; Hypocholesteremic Effect of Linoleic Acid and Phytosterol 1984, 25, CAPLUS				
(4) Hasegawa; JOSHI EIYO DAIGAKU KIYO 1983, V14, P165 CAPLUS				
(5) Kosbab, J; WO 9833494 A 1998				
ALL CITATIONS AVAILABLE IN THE RE FORMAT				

AB . . . onto the Indian market for human use. RBO contains oleic acid (38.4%), linoleic acid (34.4%), and linolenic acid (2.2%) as **unsatd. fatty acids**, and palmitic (21.5%) and stearic (2.9%) acids as satd. fatty acids. The unsaponifiable fraction (4.2%) has total tocopherols (81.3 mg%), . . . and 24-methylene cycloartanol (494 mg%). Studies on exptl. rats demonstrated a hypolipidemic effect of RBO. The unsaponifiable fraction of RBO **lowers cholesterol** levels. Feeding **phytosterols**, CA, and 24-methylene cycloartanol in amts. present in RBO to hypercholesterolemic rats for 8 wk indicates that CA alone reduces . . . and triglyceride levels significantly. Endogenous sterol excretion increases in animals given CA. The accumulation of CA in the liver inhibits **cholesterol** esterase activity, which in turn leads to **redn.** in circulating **cholesterol** levels. CA is structurally similar to cholesterol and may compete with the binding sites of cholesterol and sequester cholesterol, which. . . derivs. RBO, which is rich in tocopherols and tocotrienols, may improve oxidative stability. Tocotrienols inhibit HMG CoA reductase, resulting in **hypcholesterolemia**. The hypolipidemic effect of RBO has also been established in human subjects. Thus, RBO could be a suitable edible oil.

ACCESSION NUMBER: 1992:127428 CAPLUS
DOCUMENT NUMBER: 116:127428
TITLE: Nutritional and biochemical aspects of the hypolipidemic action of rice bran oil: a review
AUTHOR(S): Rukmini, Cheruvanky; Raghuram, Thummala C.
CORPORATE SOURCE: Natl. Inst. Nutr., Indian Counc. Med. Res., Hyderabad,
500007, India
SOURCE: J. Am. Coll. Nutr. (1991), 10(6), 593-601
CODEN: JONUDL; ISSN: 0731-5724
DOCUMENT TYPE: Journal; General Review
LANGUAGE: English
SO J. Am. Coll. Nutr. (1991), 10(6), 593-601
CODEN: JONUDL; ISSN: 0731-5724

L6 ANSWER 5 OF 11 MEDLINE

AB . . . onto the Indian market for human use. RBO contains oleic acid (38.4%), linoleic acid (34.4%), and linolenic acid (2.2%) as **unsaturated fatty acids**, and palmitic (21.5%) and stearic (2.9%) acids as saturated fatty acids. The unsaponifiable fraction (4.2%) has total tocopherols (81.3 mg%), . . . and 24-methylene cycloartanol (494 mg%). Studies on experimental rats demonstrated a hypolipidemic effect of RBO. The unsaponifiable fraction of RBO **lowers cholesterol** levels. Feeding **phytosterols**, CA, and 24-methylene cycloartanol in amounts present in RBO to hypercholesterolemic rats for 8 weeks indicates that CA alone reduces **cholesterol** and triglyceride levels significantly. Endogenous sterol excretion increases in animals given CA. The accumulation of CA in the liver inhibits **cholesterol** esterase activity, which in turn leads to **reduction** in circulating **cholesterol** levels. CA is structurally similar to **cholesterol** and may compete with the binding sites of **cholesterol** and sequester **cholesterol**, which is metabolized to its derivatives. RBO, which is rich in tocopherols and tocotrienols, may improve oxidative stability. Tocotrienols inhibit HMG CoA **reductase**, resulting in **hypcholesterolemia**. The hypolipidemic effect of RBO has also been established in human subjects. Thus, RBO could be a suitable edible oil.

ACCESSION NUMBER: 92121598 MEDLINE
DOCUMENT NUMBER: 92121598
TITLE: Nutritional and biochemical aspects of the hypolipidemic action of rice bran oil: a review.

AUTHOR: Rukmini C; Raghuram T C
CORPORATE SOURCE: National Institute of Nutrition, Indian Council of Medical Research, Hyderabad..
SOURCE: JOURNAL OF THE AMERICAN COLLEGE OF NUTRITION, (1991 Dec) 10 (6) 593-601. Ref: 36
10 Journal code: H51. ISSN: 0731-5724.
PUB. COUNTRY: United States
Journal; Article; (JOURNAL ARTICLE)
General Review; (REVIEW)
(REVIEW, TUTORIAL)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 199204
SO JOURNAL OF THE AMERICAN COLLEGE OF NUTRITION, (1991 Dec) 10 (6) 593-601.
Ref: 36
Journal code: H51. ISSN: 0731-5724.

L6 ANSWER 6 OF 11 MEDLINE
AB . . . fraction of soybean (PUFS) given over 24 weeks to 19 patients with primary types IIa and IIb hyperlipoproteinemia. The percent **reduction** of plasma **cholesterol** in types IIa and IIb were 13.2% and 11.5%, respectively. PUFS significantly reduced LDL **cholesterol** levels, but had little affect on VLDL or HDL **cholesterol**. The triglyceride/**cholesterol** ratio in HDL fraction was also significantly reduced, suggesting that PUFS plays a role in the catabolism of HDL. PUFS contained tocopherol and **unsaturated fatty acid**. The relative impact of the individual components could not be assessed directly, however, it seemed reasonable to conclude that the hypo-**cholesterolemic** effect of PUFS results from a summation effect of plant sterols, tocopherols and **unsaturated fatty acids**.

CT . . .
TU, therapeutic use
Cholesterol: BL, blood
Hypercholesterolemia, Familial: BL, blood
*Hypercholesterolemia, Familial: DT, drug therapy
Lipoproteins, LDL Cholesterol: BL, blood
***Phytosterols: TU, therapeutic use**
Soybeans
Triglycerides: BL, blood

CN 0 (Antilipemic Agents); 0 (Lipoproteins, LDL Cholesterol); 0 (**Phytosterols**); 0 (Triglycerides)

ACCESSION NUMBER: 85148557 MEDLINE

DOCUMENT NUMBER: 85148557

TITLE: Effect of the purified unsaponifiable fraction of soybean on primary type II hyperlipoproteinemia.

AUTHOR: Nakashima Y; Nakamura T; Aramaki Y; Kuroiwa A

SOURCE: ARTERY, (1983) 12 (3) 199-211.
Journal code: 8NN. ISSN: 0098-6127.

PUB. COUNTRY: United States
Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 198506

SO ARTERY, (1983) 12 (3) 199-211.
Journal code: 8NN. ISSN: 0098-6127.

L6 ANSWER 7 OF 11 EMBASE COPYRIGHT 2000 ELSEVIER SCI. B.V.

AB . . . onto the Indian market for human use. RBO contains oleic acid (38.4%), linoleic acid (34.4%), and linolenic acid (2.2%) as **unsaturated fatty acids**, and palmitic (21.5%) and stearic (2.9%) acids as saturated fatty acids. The unsaponifiable fraction (4.2%) has total tocopherols (81.3 mg%), . . . and 24-methylene

cycloartanol (494 mg%). Studies on experimental rats demonstrated a hypolipidemic effect of RBO. The unsaponifiable fraction of RBO **lowers cholesterol** levels. Feeding **phytosterols**, CA, and 24-methylene cycloartanol in amounts present in RBO to hypercholesterolemic rats for 8 weeks indicates that CA alone reduces **cholesterol** and triglyceride levels significantly. Endogenous sterol excretion increases in animals given CA. The accumulation of CA in the liver inhibits **cholesterol** esterase activity, which in turn leads to **reduction** in circulating **cholesterol** levels. CA is structurally similar to **cholesterol** and may compete with the binding sites of **cholesterol** and sequester **cholesterol**, which is metabolized to its derivatives. RBO, which is rich in tocopherols and tocotrienols, may improve oxidative stability. Tocotrienols inhibit HMG CoA **reductase**, resulting in **hypcholesterolemia**. The hypolipidemic effect of RBO has also been established in human subjects. Thus, RBO could be a suitable edible oil.

ACCESSION NUMBER: 92262702 EMBASE
DOCUMENT NUMBER: 1992262702
TITLE: Nutritional and biochemical aspects of the hypolipidemic action of rice bran oil: A review.
AUTHOR: Rukmini C.; Raghuram T.C.
CORPORATE SOURCE: National Institute of Nutrition, Hyderabad 500007, India
SOURCE: Journal of the American College of Nutrition, (1991) 10/6 (593-601).
ISSN: 0731-5724 CODEN: JONU DL
COUNTRY: United States
DOCUMENT TYPE: Journal; General Review
FILE SEGMENT: 006 Internal Medicine
017 Public Health, Social Medicine and Epidemiology
018 Cardiovascular Diseases and Cardiovascular Surgery
029 Clinical Biochemistry
LANGUAGE: English
SUMMARY LANGUAGE: English
SO Journal of the American College of Nutrition, (1991) 10/6 (593-601).
ISSN: 0731-5724 CODEN: JONU DL
L6 ANSWER 8 OF 11 WPIDS COPYRIGHT 2000 DERWENT INFORMATION LTD
TI Oil and fat composition containing **phytosterol** for preparation of food and pharmaceuticals.
AB WO 9959423 UPAB: 20000330
NOVELTY - Composition comprises a **phytosterol** dissolved in an oil and fat comprising one or more polyhydric alcohol/fatty acid esters each having an esterification degree of. . . more and containing at least one unesterified hydroxyl group.
DETAILED DESCRIPTION - An INDEPENDENT CLAIM is included for a **phytosterol** containing food which comprises one or more lipid ingredients comprising at least 15 wt.% one or more diacylglycerols. The weight ratio of diacylglycerols to **phytosterol** is 10-100.
USE - Used for preparation of oil in water type fat-processed foods such as drinks, desserts, ice creams,. . . cakes, cookies, pies, breads and chocolates, and other foods including bakery mixes, processed meat products, frozen entrees and frozen foods, **hypcholesteremic** preparation in the form of a capsule, sugar-coated tablet, molded granules, candy or drop.
ADVANTAGE - Blood cholesterol level is. . . good workability, and the flavor and feeling of food items are not changed. Generation of smoking during cooking is eliminated. **Phytosterol** is solubilized by the composition.
Dwg. 0/0
TECH. . . esterification degree of 2-4.5 and comprises glycerol/fatty acid esters, polyglycerol/fatty acid esters, sucrose/fatty acid esters or sorbitan/fatty acid esters. The **phytosterol** is contained in an

amount of 1.2 wt.% or more. 55 wt.% Fatty acids contained in diacylglycerols are **unsaturated fatty acids**.

The composition is a frying oil.

TT TT: OIL FAT COMPOSITION CONTAIN **PHYTOSTEROL** PREPARATION FOOD.
PHARMACEUTICAL.

ACCESSION NUMBER: 2000-105555 [09] WPIDS
CROSS REFERENCE: 1999-571935 [48]
DOC. NO. CPI: C2000-031612
TITLE: Oil and fat composition containing **phytosterol**
for preparation of food and pharmaceuticals.
DERWENT CLASS: A96 B01 D13
INVENTOR(S): GOTO, N; NISHIDE, T; TANAKA, Y; YASUKAWA, T
PATENT ASSIGNEE(S): (KAOS) KAO CORP
COUNTRY COUNT: 23
PATENT INFORMATION:

PATENT NO	KIND	DATE	WEEK	LA	PG
WO 9959423	A1	19991125	(200009)*	EN	26
RW: AT BE CH CY DE DK ES FI FR GB GR IE IT LU MC NL PT SE					
W: BR CA CN SG					
US 6025348	A	20000215	(200016)#		

APPLICATION DETAILS:

PATENT NO	KIND	APPLICATION	DATE
WO 9959423	A1	WO 1998-JP2228	19980521
US 6025348	A	US 1998-69754	19980430

PRIORITY APPLN. INFO: WO 1998-JP2228 19980521; US 1998-69754
19980430

L6 ANSWER 9 OF 11 WPIDS COPYRIGHT 2000 DERWENT INFORMATION LTD
TI . . . risk of cardiovascular or lipid diseases, diabetes and thyroid
disease - using as risk indicator the ratio of campesterol to beta-
sitosterol in the serum, also treatment of these disorders with
phytosterol(s).

AB WO 9801759 UPAB: 19980302
The ratio (R), in serum, of campesterol (I) to beta -**sitosterol**
(II) is determined, compared with that in a normal control and used as an
indicator of increased risk of cardiovascular. . . Also claimed are:
(1) method for assessing risk of cardiovascular and lipid disorders by
measuring (R), serum levels of total **phytosterols** (TP) and total
cholesterol (TC) and comparing all 3 values with those of normal
controls;

(2) method for increasing the inhibitory effect of **phytosterols**
(III) on enterocyte absorption of cholesterol (IV) by administering (III)
that inhibits absorption of (IV) and/or (II); (3) (IV)-lowering
composition. . . treatment of diabetes and thyroid disease or dietary
modification to increase TP and (R). (III) are hydrophobic, particularly
(I) and **sitosterol**, and may be supplied as safflower, sesame
seed, maize, rice bran, olive, rapeseed, flaxseed or coconut oils.
Particularly (III) are administered together with (i) at least 1
inhibitor

of (IV) biosynthesis, especially (A); (ii) one or more saturated or
mono/poly-**unsaturated fatty acids** (also
providing a synergistic effect) or (iii) plant oils as specified above.
Optionally the composition of (3) also includes 2-6%. . . of the (IV)
level, e.g. in subjects with apparently normal (IV) level. (III) lowers
serum TC and low-density lipoprotein (LDL) **cholesterol**, but
increases HDL and the HDL/LDL ratio, and there is a synergistic effect
when used with inhibitors (A) of 3-hydroxy-3-methylglutaryl coenzyme A
reductase, allowing a **reduction** in the dose of (A), and

thus of side effects.

Dwg.1/13

TT TT: DIAGNOSE RISK CARDIOVASCULAR LIPID DISEASE DIABETES THYROID DISEASE
RISK INDICATE RATIO BETA **SITOSTEROL** SERUM TREAT DISORDER

PHYTOSTEROL.

ACCESSION NUMBER: 1998-101200 [09] WPIDS
DOC. NO. NON-CPI: N1998-081048
DOC. NO. CPI: C1998-033490
TITLE: Diagnosing risk of cardiovascular or lipid diseases,
diabetes and thyroid disease - using as risk indicator
the ratio of campesterol to beta-**sitosterol** in
the serum, also treatment of these disorders with

phytosterol(s).

DERWENT CLASS: B04 D16 S03
INVENTOR(S): NOVAK, E
PATENT ASSIGNEE(S): (FORB-N) FORBES MEDI-TECH INC
COUNTRY COUNT: 74
PATENT INFORMATION:

PATENT NO	KIND	DATE	WEEK	LA	PG
WO 9801759	A1	19980115	(199809)*	EN	93
RW: AT BE CH DE DK ES FI FR GB GR IE IT LU MC NL PT SE					
W: AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE ES FI GB GE					
GH HU IL IS JP KE KG KP KR KZ LC LK LR LS LT LU LV MD MG MK MN MW					
MX NO NZ PL PT RO RU SD SE SG SI SK TJ TM TR TT UA UG UZ VN YU					
AU 9732518	A	19980202	(199826)		
EP 912900	A1	19990506	(199922)	EN	
R: AL AT BE CH DE DK ES FI FR GB GR IE IT LI LT LU LV MC NL PT RO SE					
SI					
US 5965449	A	19991012	(199949)		
CZ 9900011	A3	20000112	(200009)		
HU 9904042	A2	20000328	(200025)		

APPLICATION DETAILS:

PATENT NO	KIND	APPLICATION	DATE
WO 9801759	A1	WO 1997-CA474	19970703
AU 9732518	A	AU 1997-32518	19970703
EP 912900	A1	EP 1997-928091	19970703
		WO 1997-CA474	19970703
US 5965449	A	US 1996-675018	19960703
CZ 9900011	A3	WO 1997-CA474	19970703
		CZ 1999-11	19970703
HU 9904042	A2	WO 1997-CA474	19970703
		HU 1999-4042	19970703

FILING DETAILS:

PATENT NO	KIND	PATENT NO
AU 9732518	A Based on	WO 9801759
EP 912900	A1 Based on	WO 9801759
CZ 9900011	A3 Based on	WO 9801759
HU 9904042	A2 Based on	WO 9801759

PRIORITY APPLN. INFO: US 1996-675018 19960703

* L6 ANSWER 10 OF 11 BIOSIS COPYRIGHT 2000 BIOSIS
AB. . . onto the Indian market for human use. RBO contains oleic acid
(38.4%), linoleic acid (34.4%), and linolenic acid (2.2%) as
unsaturated fatty acids, and palmitic (21.5)
and stearic (2.9%) acids as saturated fatty acids. The unsaponifiable
fraction (4.2%) has total tocopherols (81.3% mg%), . . . and

24-methylene cycloartanol (494 mg%). Studies on experimental rats demonstrated a hypolipidemic effect of RBO. The unsaponifiable fraction of

RBO **lowers cholesterol** levels. Feeding **phytosterols**, CA, and 24-methylene cycloartanol in amounts present in RBO to hypercholesterolemic rats for 8 weeks indicates that CA alone reduces **cholesterol** and triglyceride levels significantly. Endogenous sterol excretion increases in animals given CA. The accumulation of CA in the liver inhibits **cholesterol** esterase activity, which in turn leads to **reduction** in circulating **cholesterol** levels. CA is structurally similar to **cholesterol** and may compete with the binding sites of **cholesterol** and sequester **cholesterol**, which is metabolized to its derivatives. RBO, which is rich in tocopherols and tocotrienols, may improve oxidative stability. Tocotrienols inhibit HMG CoA **reductase**, resulting in **hypcholesterolemia**. The hypolipidemic effect of RBO has also been established in human subjects. Thus, RBO could be a suitable edible oil. . . .

ACCESSION NUMBER: 1992:75496 BIOSIS
DOCUMENT NUMBER: BA93:43951
TITLE: NUTRITIONAL AND BIOCHEMICAL ASPECTS OF THE HYPOLIPIDEMIC ACTION OF RICE BRAN OIL A REVIEW.
AUTHOR(S): RUKMINI C; RAGHURAM T C
CORPORATE SOURCE: NATL. INST. NUTRITION, HYDERABAD 500007, INDIA.
SOURCE: J AM COLL NUTR, (1991) 10 (6), 593-601.
CODEN: JONU DL. ISSN: 0731-5724.
FILE SEGMENT: BA; OLD
LANGUAGE: English
SO J AM COLL NUTR, (1991) 10 (6), 593-601.
CODEN: JONU DL. ISSN: 0731-5724.

L6 ANSWER 11 OF 11 BIOSIS COPYRIGHT 2000 BIOSIS

AB The effects of dietary fats and **phytosterol** on the fatty acid composition and lipoprotein cholesterol in serum were studied in female rats with the following results. The. . . due to an increase in the lower density lipoprotein (LDL + VLDL [very low density lipoprotein]).

The

addition of 5% **phytosterol** to the 10% butter-cholesterol diet reduced the total cholesterol level and increased the ratio of cholesterol in high density lipoprotein. . . = 0.947), and also the level of LDL + VLDL-cholesterol (r = 0.935). Cod liver oil, wheat germ oil and **phytosterol** induce an increase in the PUFA/SFA ratio, promote **hypcholesterolemia** and change lipoprotein concentration. There were indications that no relationship exists between the change in the total cholesterol level and. . . .

IT Miscellaneous Descriptors

COD LIVER OIL WHEAT GERM OIL POLY **UNSATURATED FATTY**
-ACID SATURATED FATTY-ACID HYPO CHOLESTEROLEMIA OLEIC-ACID

ACCESSION NUMBER: 1984:349467 BIOSIS
DOCUMENT NUMBER: BA78:85947
TITLE: EFFECTS OF DIETARY FATS AND PHYTO STEROL ON SERUM FATTY-ACID COMPOSITION AND LIPO PROTEIN CHOLESTEROL IN RATS.
AUTHOR(S): HIRAI K; OHNO Y; NAKANO T; IZUTANI K
CORPORATE SOURCE: DEP. NUTR. BIOCHEM., FAC. SCI. LIVING, OSAKA CITY UNIV., SUMIYOSHI, OSAKA 558, JPN.
SOURCE: J NUTR SCI VITAMINOL, (1984) 30 (2), 101-112.
CODEN: JNSVA5. ISSN: 0301-4800.
FILE SEGMENT: BA; OLD
LANGUAGE: English
SO J NUTR SCI VITAMINOL, (1984) 30 (2), 101-112.